

Preventing Unraveling in Social Networks Gets Harder*

Rajesh Chitnis[†]

Fedor V. Fomin[‡]

Petr A. Golovach[‡]

April 25, 2013

Abstract

The behavior of users in social networks is often observed to be affected by the actions of their friends. Bhawalkar et al. [2] introduced a formal mathematical model for user engagement in social networks where each individual derives a benefit proportional to the number of its friends which are engaged. Given a threshold degree k the equilibrium for this model is a maximal subgraph whose minimum degree is $\geq k$. However the dropping out of individuals with degrees less than k might lead to a cascading effect of iterated withdrawals such that the size of equilibrium subgraph becomes very small. To overcome this some special vertices called “anchors” are introduced: these vertices need not have large degree. Bhawalkar et al. [2] considered the ANCHORED k -CORE problem: Given a graph G and integers b, k and p do there exist a set of vertices $B \subseteq H \subseteq V(G)$ such that $|B| \leq b, |H| \geq p$ and every vertex $v \in H \setminus B$ has degree at least k in the induced subgraph $G[H]$. They showed that the problem is NP-hard for $k \geq 2$ and gave some inapproximability and fixed-parameter intractability results. In this paper we give improved hardness results for this problem. In particular we show that the ANCHORED k -CORE problem is W[1]-hard parameterized by p , even for $k = 3$. This improves the result of Bhawalkar et al. [2] (who show W[2]-hardness parameterized by b) as our parameter is always bigger since $p \geq b$. Then we answer a question of Bhawalkar et al. [2] by showing that the ANCHORED k -CORE problem remains NP-hard on planar graphs for all $k \geq 3$, even if the maximum degree of the graph is $k + 2$. Finally we show that the problem is FPT on planar graphs parameterized by b for all $k \geq 7$.

1 Introduction

A social network can be thought as the graph of relationships and interactions within a group of individuals. Social networks play a leading role in various fields such as social sciences [11, 12], life sciences [6, 20] and medicine [6, 15]. Social networks today perform a fundamental role as a medium for the spread of information, ideas, and influence among its members. As an example, Facebook reported a figure of one billion active users as of October 2012 [1]. An important characteristic of social networks is that the behavior of an individual is often influenced by the actions of their friends. New events occur quite often in social networks: some examples are usage of a particular cell phone brand, adoption of a new drug within the medical profession, or the rise of a political movement in an unstable society. To estimate whether these events or ideas spread extensively or die out soon, we need to model and study the dynamics of *influence propagation* in social networks. We consider the following model of *user engagement* defined by Bhawalkar

*Supported by the European Research Council (ERC) via grant Rigorous Theory of Preprocessing, reference 267959

[†]Dept. of Computer Science, University of Maryland at College Park, USA. Supported in part by NSF CAREER award 1053605, NSF grant CCF-1161626, ONR YIP award N000141110662, DARPA/AFOSR grant FA9550-12-1-0423, a University of Maryland Research and Scholarship Award (RASA) and a Summer International Research Fellowship from the University of Maryland. Email: rchitnis@cs.umd.edu

[‡]Department of Informatics, University of Bergen, PB 7803, 5020 Bergen, Norway. Email: {fedor.fomin, petr.golovach}@ii.uib.no

et al. [2]: there is a single product and each individual has two options of “engaged” or “drop out”. Initially we assume that all individuals are engaged. There is a given threshold parameter k such that a person finds it worthwhile to remain engaged if and only if at least k of her friends are still engaged. For example engagement could represent active participation in a social network, and individuals might drop out and switch to a new social network if less than k of his friends are active on the current social network. Indeed such a phase transition has been observed in the popularity of social networks: in India the leading social network was Orkut until Facebook surpassed it in August 2010 [9].

In our model of *user engagement* all individuals with less than k friends will clearly drop out. Unfortunately this can be contagious and may affect even those individuals who initially had more than k friends in the social network. An extreme example of this is given in page 17 of [17]: consider a path on n vertices and let $k = 2$. Note that $n - 2$ vertices have degree two in the network. However there will be a *cascade of iterated withdrawals*. An endpoint has degree one, it drops out and now its neighbor in the path has only one friend in the social network and it drops out as well. It is not hard to see that the whole network eventually drops out. In general at the end of all the iterated withdrawals the remaining engaged individuals form a unique maximal induced subgraph whose minimum degree is at least k . This is called as the *k-core* and is a well-known concept in the theory of social networks. It was introduced by Seidman [19] and also been studied in various social sciences literature [3, 4].

A Game-Theoretic Model: Consider the following game-theoretical model [2]: each user in a social network pays a cost of k to remain engaged. On the other hand it receives a profit of 1 from every neighbor who is engaged. The “network effects” come into play, and an individual decides to remain engaged if has non-negative payoff, i.e., it has at least k neighbors who are engaged. The *k-core* can be viewed as the unique maximal equilibrium in this model. Assuming that all the players make decisions simultaneously the model can be viewed as a simultaneous-move game where each individual has two strategies viz. remaining engaged or dropping out. Consider a graph G defined on the set of players by adding an edge between two players if and only if they are friends in the network. For every strategy profile δ let S_δ denote the set of players who remain engaged. The payoff for a person v is 0 if she is not engaged, otherwise it is the number of her friends among engaged players minus k . We can easily characterize the set of pure Nash equilibria for this game: a strategy profile δ is a Nash equilibrium if and only if the following two conditions hold:

- No engaged player wants to drop out, i.e., minimum degree of the induced graph $G[S_\delta]$ is $\geq k$
- No player who has dropped out wants to become engaged, i.e., no $v \in V(G) \setminus S_\delta$ has $\geq k$ neighbors in S_δ

In general there can be many Nash equilibria. For example, if G itself has minimum degree $\geq k$ then $S_\delta = \emptyset$ and $S_\delta = V(G)$ are two equilibria (and there may be more). Recall that the goal of the product company is to attract as many people as possible. Owing to the fact that it is a maximal equilibrium, the *k-core* has the special property that it is beneficial to both parties: it maximizes the payoff of every user, while also maximizing the payoff of the product company. Chwe [3, 4] and Sääskilahti [16] claim that one can reasonably expect this maximal equilibrium even in real-life implementations of this game.

Preventing Unraveling: The unraveling described above in Schelling’s example of a path is highly undesirable since the goal is to keep as many people engaged as possible. How can we attempt to prevent this unraveling? In Schelling’s example it is easy to see: if we “buy” the two end-point players into being engaged then the whole path becomes engaged.

In general we overcome the issue of unraveling by allowing some “anchors”: these are vertices that remain engaged irrespective of their payoffs. This can be achieved by giving them extra incentives or discounts. The hope is that with a few *anchors* we can now ensure a large subgraph remains engaged. This subgraph is now called as the *anchored k-core*: each non-anchor vertex in this induced subgraph must

have degree at least k while the anchored vertices can have arbitrary degrees. We use the notation $\deg_S(v)$ to denote the degree of v in the graph S . Bhawalkar et al. [2] formally defined the ANCHORED k -CORE problem :

The ANCHORED k -CORE Problem (AKC)

Input : An undirected graph $G = (V, E)$ and integers b, k

Question: Find a set of vertices $H \subseteq V$ of maximum size such that

- There is a set $B \subseteq H$ and $|B| \leq b$
- Every $v \in H \setminus B$ satisfies $\deg_{G[H]}(v) \geq k$

The AKC problem deals with finding a small group of individuals whose engagement is essential for the health of the social network. We call the set B as *anchors*, the set H as the *anchored k -core* and the set $H \setminus B$ as the *anchored k -supercore*. The decision version of the ANCHORED k -CORE problem deals with anchoring a given number of vertices to maximize the number of engaged vertices. More formally:

p -AKC

Input : An undirected graph $G = (V, E)$ and integers b, k, p

Question: Do there exist sets $B \subseteq H \subseteq V$ such that

- $|B| \leq b$ and $|H| \geq p$
- Every $v \in H \setminus B$ satisfies $\deg_{G[H]}(v) \geq k$

Previous Work: Bhawalkar et al. [2] introduced the ANCHORED k -CORE problem and gave some positive and negative results for this problem. Noting that the problem is trivial for $k = 1$, they showed that AKC is polynomial time solvable for $k = 2$ but NP-hard for all $k \geq 3$. They also gave a strong inapproximability result: it is NP-hard to approximate the AKC problem to within an $O(n^{1-\epsilon})$ factor for any $\epsilon > 0$. From the viewpoint of parameterized complexity they showed that for every $k \geq 3$ the p -AKC problem is W[2]-hard with respect to b . Finally on the positive side they give a polynomial time algorithm on graphs of bounded treewidth. On graphs with treewidth at most w their algorithm runs in $O(3^w(k+1)^{2w}b^2) \cdot \text{poly}(n)$ time.

Our Results: It is easy to see that ANCHORED k -CORE can be solved in time $n^{b+O(1)}$, as we can try all subsets B of size b of the set of vertices of the input graph, and for each B , find the unique k -core H of maximum size such that $\deg_{G[H]}(v) \geq k$ if $v \in H \setminus B$ by the consecutive deletions of small degree vertices. We show that this result is optimal in some sense by proving that p -AKC problem is W[1]-hard parameterized by $b + k + p$. We also show that the p -AKC problem is W[1]-hard parameterized by p even for $k = 3$. This improves the result of Bhawalkar et al. [2] (who show W[2]-hardness parameterized by b), because our parameter is always bigger since $p \geq b$. Bhawalkar et al. raised the question of resolving the complexity of the ANCHORED k -CORE problem on special graph classes. In this paper we consider the complexity of the AKC problem on the class of planar graphs. We show that the ANCHORED k -CORE problem is NP-hard on planar graphs for all $k \geq 3$, even if the maximum degree of the graph is $k + 2$. Finally on the positive side we show that the p -AKC problem on planar graphs is FPT parameterized by b for all $k \geq 7$.

2 Fixed-Parameter Intractability Results

In this section we give two parameterized intractability results. Before that we give a brief introduction to parameterized complexity.

Parameterized Complexity: *Parameterized Complexity* is basically a two-dimensional generalization of “P vs. NP” where in addition to the overall input size n , one studies the effects on computational complexity of a secondary measurement that captures additional relevant information. This additional information can be, for example, a structural restriction on the input distribution considered, such as a bound on the treewidth of an input graph or the size of solution set. For general background on the theory see [7, 10, 14]. For decision problems with input size n , and a parameter k , the two dimensional analogue (or generalization) of P, is solvability within a time bound of $O(f(k)n^{O(1)})$, where f is a computable function of k alone. Problems having such an algorithm are said to be *fixed parameter tractable* (FPT). Such algorithms are practical when small parameters cover practical ranges. The W -hierarchy is a collection of computational complexity classes: we omit the technical definitions here. The following relation is known amongst the classes in the W -hierarchy: $FPT = W[0] \subseteq W[1] \subseteq W[2] \subseteq \dots$. It is widely believed that $FPT \neq W[1]$, and hence if a problem is hard for the class $W[i]$ (for any $i \geq 1$) then it is considered to be fixed-parameter intractable.

W[1]-hardness parameterized by $b+k+p$: In this section we show that the p -AKC problem is $W[1]$ -hard even when parameterized by $b+k+p$. We reduce from the well-known $W[1]$ -hard problem CLIQUE

CLIQUE

Input : An undirected graph $G = (V, E)$ and an integer ℓ

Question: Does G have a clique of size at least ℓ ?

Theorem 1. *The p -AKC problem is $W[1]$ -hard parameterized by $b+k+p$ for $k \geq 3$.*

Proof. Consider an instance $G = (V, E)$ of CLIQUE. Let $V = \{v_1, v_2, \dots, v_n\}$. For each edge $e = \{v_i, v_j\}$ we subdivide it and add a new vertex $w_{\{i,j\}}$. Let this new graph be $G' = (V', E')$. Define $b = \binom{\ell}{2}$, $k = \ell - 1$ and $p = \ell + \binom{\ell}{2}$. The claim is the instance (G', b, k, p) of p -AKC answers YES and only if the instance (G, ℓ) of CLIQUE answers YES.

Suppose G has a clique of size ℓ say $C = \{v_1, v_2, \dots, v_\ell\}$. Pick as anchors in G' the $\binom{\ell}{2}$ vertices of the type $w_{\{i,j\}}$ for $1 \leq i \neq j \leq \ell$. It is easy to see that these anchors along with the vertex set C form a k -core of size $\ell + \binom{\ell}{2}$ for the instance (G', b, k, p) of p -AKC.

Suppose that the instance (G', b, k, p) of p -AKC has a solution. Since any newly added vertex has degree two we know that every vertex in the k -supercore must be from V . Further each vertex from the supercore needs at least k anchors, and each anchor can be shared between at most two vertices from k -supercore. Let S denote the k -supercore. Counting the number of anchors (with repetitions) in two ways we have $2\binom{\ell}{2} \geq |S|(\ell - 1)$, i.e., $\ell \geq |S|$. But we know that $|S| \geq p - b \geq \ell$, and hence $|S| = \ell$. Without loss of generality let the vertices of S be v_1, v_2, \dots, v_ℓ . Each v_i has degree at least $k = \ell - 1$ in $H = S \cup B$, and so S must form a clique in G . \square

3 W[1]-hardness parameterized by p

Bhawalkar et al. (Theorem 3 in [2]) showed that the p -AKC problem is $W[2]$ -hard parameterized by b for every $k \geq 3$. In this section we prove that it is in fact $W[1]$ -hard parameterized by p for $k = 3$. This improves on the result of Bhawalkar et al. [2] in two aspects: firstly our parameter is stronger since $p \geq b$ by definition. Also we show hardness for a smaller complexity class since it is known that $FPT \subseteq W[1] \subseteq W[2]$.

Theorem 2. *The p -AKC problem is $W[1]$ -hard parameterized by p for $k = 3$.*

Proof. We again reduce from the CLIQUE problem. Consider an instance $(G = (V, E), \ell)$ of CLIQUE where $V = (v^1, v^2, \dots, v^n)$. Construct a new graph $G' = (V', E')$ as follows. For each $1 \leq i \neq j \leq \ell$ make a copy G_{ij} of the vertex set V (do not add any edges). Let the vertex v^r in the copy G_{ij} be labeled v_{ij}^r . Add the following edges to G' :




Figure 1: The graph G' constructed in Theorem 2 for the special case when $n = 8$ and $\ell = 5$.

- For each $1 \leq i \neq j \leq \ell$ and $r, s \in [\ell]$ we add an edge between v_{ij}^r and v_{ji}^s if and only if $v^r v^s \in E$. Subdivide each such edge by adding a new vertex called *green*.
- For each $1 \leq i, j \leq \ell$ add the cycle $v_{i1}^j - v_{i2}^j - \dots - v_{i,i-1}^j - v_{i,i+1}^j - \dots - v_{i\ell}^j - v_{i1}^j$. Let us denote this cycle by C_i^j .

This completes the construction of G' . Let $k = 3$, $b = \binom{\ell}{2}$ and $p = 3b$. The claim is that the instance (G, ℓ) of CLIQUE answers YES if and only if the instance (G', b, k, p) of p -AKC answers YES.

Suppose G has a clique of size ℓ , say $C = \{v_1, v_2, \dots, v_\ell\}$. For $1 \leq i \neq j \leq \ell$ pick the vertex v_{ij}^i from G_{ij} . This gives a set S of $2b$ vertices. It is easy to see that $G'[S]$ consists of disjoint cycles, and hence is regular of degree two. Now for every v_{ij}^i there is a unique vertex v_{ji}^j which is connected to it by a subdivided edge in G' . The green vertices of this subdivided edges become the anchors. Note that we pick exactly $\frac{|S|}{2} = b$ anchors. It is easy to see that each vertex of S now has degree three in the resulting induced subgraph, and hence S becomes the k -supercore. Therefore the instance (G', b, k, p) of p -AKC answers YES.

Now suppose that the instance (G', b, k, p) of p -AKC answers YES. Let S be the k -supercore and B be the set of anchors. Then we know that $|S| \geq p - b = 2b$. Each green vertex has degree two in G' , and hence cannot be in S . Each vertex in S needs at least one green vertex to achieve degree three in $G'[S \cup B]$. But any green vertex can be used by at most two vertices in S . Therefore we have $2b \geq 2|B| \geq |S| \geq 2b$ which implies $|B| = b$ and $|S| = 2b$. Hence the budget must come from the green vertices only, and that the vertices of S form a *matching* under the relation of sharing a common green vertex. Without loss of generality let $v_{12}^{i_1}$ and $v_{21}^{i_2}$ be two vertices in S such that they share a green vertex from B . Now we know that $v_{12}^{i_1}$ has degree at least three in $G'[B \cup S]$ but cannot be incident to any other green vertex. So we need to include $v_{13}^{i_1}$ and $v_{1\ell}^{i_1}$ in S . Again each of these two vertices can be incident to at most one green vertex in $G'[B \cup S]$ and ultimately this means that we must have $C_1^{i_1} \subseteq G'[B \cup S]$. For $2 \leq j \leq \ell$ we know that the vertex $v_{1j}^{i_1}$ needs one more

edge to achieve degree at least three. This edge must be towards a green vertex which is adjacent to some vertex in G_{j1} , say v_{j1}^{ij} . By reasoning similar to above we must have $C_j^{ij} \subseteq G'[B \cup S]$ for every $2 \leq j \leq \ell$. So we have chosen $2b$ vertices in S , which is the maximum allowed budget. Therefore $S \cap G_{jj'} = \{v_{jj'}^{ij}\}$ for every $1 \leq j \neq j' \leq \ell$.

The claim is that the set $\{v^{i1}, v^{i2}, \dots, v^{i\ell}\}$ forms a clique in G . Consider any two indices $1 \leq q \neq r \leq \ell$. We know that the vertex v_{qr}^{iq} is in S and has degree two in $G'[B \cup S]$ as it is in the cycle C_q^{iq} . To achieve degree three it must be incident to some green vertex. Also it must share this green vertex with some other vertex from $G_{rq} \cap S$. But we know that $G_{rq} \cap S = \{v_{rq}^{ir}\}$. Therefore v_{qr}^{iq} and v_{rq}^{ir} share a green vertex, i.e., v^{iq} and v^{ir} are adjacent in G , i.e., the vertices $\{v^{i1}, v^{i2}, \dots, v^{i\ell}\}$ form a clique in G . \square

4 NP-hardness Results on Planar Graphs

Bhawalkar et al. [2] raised the question of investigating the complexity of the ANCHORED k -CORE problem on special cases of graphs such as planar graphs. In this section we provide some answers by showing NP-hardness results for planar graphs for $k \geq 3$. The case $k \geq 4$ can be handled by a single reduction, but $k = 3$ is more complicated and requires a separate reduction. We reduce from the following problem which was shown to be NP-hard by Dahlhaus et al. [5]:

RESTRICTED-PLANAR-3-SAT

Input : A Boolean CNF formula ϕ such that

- Each clause has at most 3 literals
- Each variable is used in at most 3 clauses
- Each variable is used at least once in positive and at least once in negation
- The graph G_ϕ (described below) is planar

Question: Is the formula ϕ satisfiable ?

Consider an instance ϕ of RESTRICTED-PLANAR-3-SAT with variables x_1, x_2, \dots, x_n and clauses C_1, C_2, \dots, C_m . We associate the following graph G_ϕ with ϕ :

- For each $1 \leq i \leq n$ introduce the vertices r_i, x_i and \bar{x}_i . Add the edges $r_i x_i$ and $r_i \bar{x}_i$.
- For each $1 \leq j \leq m$ introduce the vertex c_j .
- For each $1 \leq i \leq n$ and $1 \leq j \leq m$ add an edge between x_i (or \bar{x}_i) and c_j iff x_i (or \bar{x}_i) belongs to the clause C_j .

4.1 NP-hardness on Planar Graphs for $k = 3$

In this section we show that the ANCHORED k -CORE problem is NP-hard on planar graphs for all $k = 3$, even in graphs of maximum degree 5.

Theorem 3. *For $k = 3$ the ANCHORED k -CORE problem is NP-hard even on planar graphs of maximum degree 5.*

Proof. We reduce from the RESTRICTED-PLANAR-3-SAT problem. For an instance ϕ of RESTRICTED-PLANAR-3-SAT let G_ϕ be the associated planar graph. We define two special graphs T and W (see Figure 2) and build the graph G as follows (see Figure 3):

- For each $1 \leq i \leq n$ subdivide the edge $r_i x_i$ and let the newly introduced vertex be y_i




Figure 2: The graphs T and W used in construction of G in Theorem 3. Note that T has $2n + 3$ vertices, and exactly one vertex has degree two. The graph W has exactly one vertex of degree one.

- For each $1 \leq i \leq n$ subdivide the edge $r_i \bar{x}_i$ and let the newly introduced vertex be \bar{y}_i .
- For each $1 \leq i \leq n$ attach a copy of T by identifying its degree two vertex with vertex r_i . Call this gadget as R_i
- For each $1 \leq i \leq n$ if x_i appears in exactly one clause then attach a copy of W by identifying its degree one vertex with vertex x_i . Call this gadget as X_i
- For each $1 \leq i \leq n$ if \bar{x}_i appears in exactly one clause then attach a copy of W by identifying its degree one vertex with vertex \bar{x}_i . Call this gadget as \bar{X}_i
- For each $1 \leq j \leq m$ attach a copy of T by identifying its degree two vertex with vertex c_j . Call this gadget as C_j

This completes the construction of the graph $G = (V, E)$. All the gadgets we added are planar, and it is easy to verify that the planarity is preserved when we construct G from G_ϕ . Let $k = 3$, $b = n$ and $p = |V| - 2n$. The claim is that ϕ is satisfiable if and only if the instance (G, b, k, p) of p -AKC answers YES.

Suppose ϕ is satisfiable. For each $1 \leq i \leq n$, if $x_i = 1$ in the satisfying assignment for ϕ then pick y_i in B and \bar{x}_i, \bar{y}_i in B' . Otherwise pick \bar{y}_i in B and x_i, y_i in B' . Clearly $|B| = n$ and $|B'| = 2n$. Let B be the set of anchors and set $H = V \setminus B'$. Now the claim is that every vertex $w \in H \setminus B$ has degree at least three in the induced subgraph $G[H]$. For each $1 \leq i \leq n$, exactly one of y_i or \bar{y}_i is in B . Hence r_i (and also each vertex of R_i) has degree exactly three in H . Consider a literal x_i . We have the following two cases:

- $y_i \in B$: x_i gets one edge from y_i . If x_i appears in exactly one clause then it gets one edge from that clause vertex and one edge from its neighbor in X_i (and each vertex in X_i has degree at least three in H). Otherwise x_i gets two edges from the two clause vertices which it appears in.
- $\bar{y}_i \in B$: \bar{x}_i gets one edge from \bar{y}_i . If \bar{x}_i appears in exactly one clause then it gets one edge from that clause vertex and one edge from its neighbor in \bar{X}_i (and each vertex in \bar{X}_i has degree at least three in H). Otherwise \bar{x}_i gets two edges from the two clause vertices which it appears in.

Finally consider a clause vertex c_j . It has at least one true literal say x_i in it. In addition c_j has two neighbors in C_j , and hence the degree of c_j is at least three in H . Consequently each vertex in C_j has degree at least three in H . Therefore with $b = |B| = n$ anchors we can cover a 3-core of size at least $|V \setminus B'| = |V| - 2n = p$, and hence (G, b, k, p) answers YES.

Suppose that the instance (G, b, k, p) of p -AKC answers YES. Let us denote the 3-core by H . Note that we can afford to not have at most $2n$ vertices in the 3-core. Each y_i and \bar{y}_i have degree two in G : so either




Figure 4: The graph G constructed in Theorem 4 for $k = 4$.

1. For each $1 \leq i \leq n$

- Add a set Y_i of $k - 1$ vertices and make all of them adjacent to r_i .
- For each vertex $y \in Y_i$ add $k - 1$ vertices and make all of them adjacent to y . Let Z_i be the set of all these $(k - 1)^2$ vertices.

2. For each $1 \leq j \leq m$

- Add a set U_j of $k - 1$ vertices and make all of them adjacent to v_j .
- For each vertex $u \in U_j$ add $k - 1$ vertices and make all of them adjacent to u . Let W_j be the set of all these $(k - 1)^2$ vertices.

Set $b = n((k - 1)^2 + 1) + m(k - 1)^2$ and $p = n(k(k - 1) + 2) + m(k(k - 1) + 1) = b + nk + mk$. Note that degree of each x_i and \bar{x}_i is at most three in G . We claim that ϕ is satisfiable if and only if the instance (G, b, k, p) of p -AKC answers YES.

Suppose ϕ is satisfiable. For each $1 \leq i \leq n$, if $x_i = 1$ in the satisfying assignment then select x_i in B' and \bar{x}_i in B'' . Else select \bar{x}_i in B' and x_i in B'' . Let $B = B' \cup (\cup_{i=1}^n Z_i) \cup (\cup_{j=1}^m W_j)$. Let us place the anchors at vertices of B . Then $|B| = n + n(k - 1)^2 + m(k - 1)^2 = b$. Now the claim is that $V \setminus B''$ forms a k -core. This would conclude the proof since $|V \setminus B''| = |G| - n = p$. For each $1 \leq i \leq n$ the vertex r_i gets one neighbor from either x_i or \bar{x}_i and it has $k - 1$ neighbors in Y_i . Each vertex in Y_i has one neighbor in r_i and $k - 1$ neighbors in Z_i . Each vertex in U_j has one neighbor in v_j and $k - 1$ neighbors in W_j . Since there is a satisfying assignment we know that v_j has at least one neighbor in B' , and of course has $k - 1$ neighbors in U_j . So $V \setminus B''$ forms a k -core with B as the anchor set, and the instance (G, b, k, p) of p -AKC answers YES.

Now suppose that the instance (G, b, k, p) of p -AKC answers YES. Let us denote the anchors by B and the k -core by H . Consider $S = (\cup_{i=1}^n \{x_i, \bar{x}_i\}) \cup (\cup_{i=1}^n Z_i) \cup (\cup_{j=1}^m W_j)$. Note that $|S| = b + n$. Any vertex in S has degree at most $\max\{k - 1, 3\}$ in G : so if it is present in the k -core then it must be an anchor. Since $p = |V| - n$ at least $|S| - n$ vertices from S must be anchors. Since $|S| - n = b$ these vertices must be the anchor set say B and the k -core is $H = B \cup (V \setminus S)$. Let $z \in Z_i$ for some $1 \leq i \leq n$ be adjacent to $y \in Y_i$ in

G . If $z \notin B$ then y has at most $k - 1$ neighbors in H , which contradicts the fact that $Y_i \subseteq H \setminus B$. Therefore $Z_i \subseteq B$ for every $1 \leq i \leq [n]$. Similarly we have $W_j \subseteq B$ for every $1 \leq j \leq m$. So now we can only choose n more anchors from the set $\cup_{i=1}^n \{x_i, \bar{x}_i\}$. Suppose for some $1 \leq i \leq n$ we have both $x_i \notin B$ and $\bar{x}_i \notin B$. Then the vertex r_i has degree at most $k - 1$ in $G[H]$, contradicting the fact that $r_i \in H \setminus B$. Therefore for every $1 \leq i \leq n$ at least one of x_i or \bar{x}_i must be in B . As the budget for the anchors is n we know that for every $1 \leq i \leq n$ exactly one of x_i or \bar{x}_i is in B . Consider the assignment $f : \{1, 2, \dots, n\} \rightarrow \{0, 1\}$ for ϕ given by $f(i) = 1$ if $x_i \in B$ and 0 otherwise. The claim is that f is a satisfying assignment for ϕ . Consider any clause C_j of ϕ . The vertex v_j has exactly $k - 1$ neighbors in U_j , and hence must have at least one neighbor in B (which appears in the clause C_j). If this neighbor is some $x_i \in B$ then the assignment would set $x_i = 1$. Else the neighbor is of the type $\bar{x}_i \in B$ and then our assignment would have set $x_i = 0$. Hence f is a satisfying assignment for ϕ . Finally note that the maximum degree of G is $k + 2$, which can occur if v_j has three literals. \square

5 FPT on Planar Graphs Parameterized by b

In this section we show that the p -AKC problem is FPT parameterized by b on planar graphs when $k \geq 7$.

Lemma 1. *The problem of checking whether there is an anchored k -core such that $q \geq |H| \geq p$ can be expressed in first-order logic.*

Proof. Consider the following formula in first-order logic:

$$\phi_q = \bigvee_{p \leq i \leq q} (\exists h_1, h_2, \dots, h_i : H_i \wedge \bigvee_{1 \leq j \leq b} (\exists b_1, b_2, \dots, b_j : B_j \wedge B_j H_i \wedge \forall y : (\bigvee_{1 \leq i_1 \leq i} (y = h_{i_1}) \wedge Y B_j) \rightarrow \exists v_1 \dots v_k : V_k \wedge V_k H_i \wedge Y V_k))$$

where

$$H_i = \bigwedge_{1 \leq i_1 \neq i_2 \leq i} (h_{i_1} \neq h_{i_2})$$

$$B_j = \bigwedge_{1 \leq j_1 \neq j_2 \leq j} (b_{j_1} \neq b_{j_2})$$

$$B_j H_i = \bigwedge_{1 \leq j_1 \leq j} (\bigvee_{1 \leq i_1 \leq i} (b_{j_1} = h_{i_1}))$$

$$Y B_j = \bigwedge_{1 \leq j_1 \leq j} (y \neq b_{j_1})$$

$$V_k = \bigwedge_{1 \leq k_1 \neq k_2 \leq k} (v_{k_1} \neq v_{k_2})$$

$$V_k H_i = \bigwedge_{1 \leq k_1 \leq k} (\bigvee_{1 \leq i_1 \leq i} (v_{k_1} = h_{i_1}))$$

$$Y V_k = \bigwedge_{1 \leq k_1 \leq k} (y v_{k_1} \in E)$$

We claim that the formula ϕ_q correctly expresses the problem of checking whether there is an anchored k -core such that $q \geq |H| \geq p$. The formulae H_i , B_j and B_k just check that all the variables in the respective formula are pairwise distinct. The formula $B_j H_i$ checks every anchor is present in the anchored k -core H . Finally for every $y \in H \setminus B$ we enforce that there are at least k elements v_1, v_2, \dots, v_k which are pairwise distinct, present in H and adjacent to y . It is now easy to see that any solution H such that $q \geq |H| \geq p$ gives a solution to the formula ϕ_q and vice versa, i.e., the formula ϕ_q exactly expresses this problem. Note that the length of ϕ_q is $\text{poly}(q)$ since $q \geq p \geq b$ and $q \geq k - 1$. \square

Seese [18] showed that any graph problem expressible in first-order logic can be solved in linear FPT time on graphs of bounded degree. More formally, let \mathcal{X} be a graph problem and $\phi_{\mathcal{X}}$ be a first-order formula for \mathcal{X} . For a constant $c > 0$ consider the graph class $\mathcal{G}_c = \{ G \mid \Delta(G) \leq c \}$. Then for every $G \in \mathcal{G}$ we can solve \mathcal{X} in $O(f(|\phi_{\mathcal{X}}|) \cdot |G|)$ where f is some function. This was later extended by Dvorak et al. [8] to a much richer graph class known as graphs with *bounded expansion*. We refer to [8, 13] for the exact definitions. However we remark that examples of such graph classes are graphs of bounded degree, graphs of bounded genus (including planar graphs), graphs that exclude a fixed (topological) minor, etc. Using these results we can give FPT algorithm parameterized by b on some classes of sparse graphs when k is sufficiently large. However for the sake of simplicity we just state the result for planar graphs.

Lemma 2. *Let G be a planar graph on n vertices. Let $k \geq 7$ and m be the set of vertices of degree at least k . Then $\frac{m}{n} < \frac{6}{7}$.*

Proof. Summing up the degrees of the graph gives twice the number of edges. For $v \in G$ we define $\deg'(v) = 7$ if $\deg(v) \geq 7$, and $\deg'(v) = 1$ otherwise. Note that every $v \in G$ satisfies $\deg'(v) \leq \deg(v)$. It is a well known fact that the maximum number of edges in a n vertex planar graph is $3n - 6$. Therefore we have $2(3n - 6) = 6n - 12 \geq 2|E(G)| = \sum_{v \in G} \deg(v) \geq \sum_{v \in G} \deg'(v) = 7m + (n - m)$. Rearranging we get $6m \leq 5n - 12$ which implies $\frac{m}{n} \leq \frac{5n-12}{6n} < \frac{6}{7}$. \square

Now we show that for $k \geq 7$ the p -AKC problem is FPT parameterized by b on planar graphs.

Theorem 5. *Let $k \geq 7$. Then for the class of planar graphs the p -AKC problem can be solved in linear FPT time parameterized by the number of anchors b .*

Proof. Fix any $k \geq 7$. Suppose there is a solution H for the p -AKC problem and let B be the set of anchors. Each vertex in $H \setminus B$ has degree at least k . Applying Lemma 2 to the planar graph $G[H]$ we have $\frac{|H| - |B|}{|H|} = \frac{|H \setminus B|}{|H|} < \frac{6}{7}$. Therefore $\frac{|B|}{|H|} \geq \frac{1}{7}$ and so $7b \geq 7|B| \geq |H|$. So we can express the problem as checking whether there is an anchored k -core H such that $p \leq |H| \leq 7b$. As shown in Lemma 1 we write the first-order formula ϕ_{7b} for this problem. By the result of Dvorak et al. [8] the p -AKC problem can be solved in $O(f(7b) \cdot n)$ for some function f , i.e., it can be solved in linear FPT time parameterized by b . \square

6 Conclusions and Open Problems

We studied the complexity of the AKC problem on the class of planar graphs, thus answering the question raised in [2]. We showed that the AKC problem is NP-hard on planar graphs, even if the graph has maximum degree $k + 2$. We also improve some fixed-parameter intractability results for the p -AKC problem. Finally on the positive side we show that for all $k \geq 7$ the p -AKC problem on planar graphs is FPT parameterized by b .

There are still several interesting questions remaining. We mention some of them here: what is the parameterized complexity status of the problem parameterized by b on planar graphs for $3 \leq k \leq 6$? What happens when we consider the problem on random graphs? Can we get reasonable approximation algorithms on some restricted graph classes?

Acknowledgements: We would like to thank the anonymous referees of AAAI 2013 for helpful comments.

References

- [1] BBC. Facebook surpasses one billion users as it tempts new markets. 2012.
- [2] K. Bhawalkar, J. M. Kleinberg, K. Lewi, T. Roughgarden, and A. Sharma. Preventing Unraveling in Social Networks: The Anchored k -Core Problem. In *ICALP* (2), pages 440–451, 2012.
- [3] M. Chwe. Structure and Strategy in Collective Action 1. *American Journal of Sociology*, 105(1):128–156, 1999.
- [4] M. Chwe. Communication and Coordination in Social Networks. *The Review of Economic Studies*, 67(1):1–16, 2000.

- [5] E. Dahlhaus, D. S. Johnson, C. H. Papadimitriou, P. D. Seymour, and M. Yannakakis. The Complexity of Multiterminal Cuts. *SIAM J. Comput.*, 23(4):864–894, 1994.
- [6] Z. Dezső and A. Barabási. Halting Viruses in Scale-free Networks. *Physical Review E*, 65(5):055103, 2002.
- [7] R. G. Downey and M. R. Fellows. *Parameterized Complexity*. Springer-Verlag, New York, 1999.
- [8] Z. Dvorak, D. Král, and R. Thomas. Deciding First-Order Properties for Sparse Graphs. In *FOCS*, pages 133–142, 2010.
- [9] Facebook. Google Makes Change to Orkut as Facebook Wins in India (<http://blogs.wsj.com/digits/2010/08/24/google-makes-change-to-orkut-as-facebook-wins-in-india/>). 2005.
- [10] J. Flum and M. Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. An EATCS Series. Springer-Verlag, Berlin, 2006.
- [11] R. T. Mikolajczyk and M. Kretzschmar. Collecting social contact data in the context of disease transmission: Prospective and retrospective study designs. *Social Networks*, 30(2):127 – 135, 2008.
- [12] S. Milgram. The Small World Problem. *Psychology Today*, 2:60–67, 1967.
- [13] J. Nešetřil and P. O. de Mendez. Linear time low tree-width partitions and algorithmic consequences. In *STOC*, pages 391–400, 2006.
- [14] R. Niedermeier. *Invitation to fixed-parameter algorithms*, volume 31 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2006.
- [15] R. Pastor-Satorras and A. Vespignani. Epidemic Spreading in Scale-free Networks. *Physical review letters*, 86(14):3200–3203, 2001.
- [16] P. Sääskilähti. Monopoly Pricing of Social Goods. *Technical Report. University Librray of Munich*, 2007.
- [17] T. Schelling. *Micromotives and Macrobbehavior*. WW Norton, 2006.
- [18] D. Seese. Linear Time Computable Problems and First-Order Descriptions. *Mathematical Structures in Computer Science*, 6(6):505–526, 1996.
- [19] S. Seidman. Network Structure and Minimum Degree. *Social networks*, 5(3):269–287, 1983.
- [20] D. Wilson. Levels of Selection: An Alternative to Individualism in Biology and the Human Sciences. *Social Networks*, 11(3):257–292, 1989.